# This Page Is Inserted by IFW Operations and is not a part of the Official Record

# **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

# IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

## WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



(51) International Patent Classification <sup>5</sup> :		(11) International Publication Number: WO 94/12599
C11D 1/66, 1/72, 1/722, 1/825	A1	(43) International Publication Date: 9 June 1994 (09.06.94)
(21) International Application Number: PCT/US9		9, B-1933 Sterrebeek (BE).
(22) International Filing Date: 19 November 1993 (1	9.11.9	(74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45202 (US).
(30) Priority Data: 92870191.1 26 November 1992 (26.11.92	2) E	
(34) Countries for which the regional or	•	
international application was filed: I 93870050.7 19 March 1993 (19.03.93)	BE et a E	.   ( - ,
(34) Countries for which the regional or		RO, RU, SD, SK, UA, US, UZ, VN, OAPI patent (BF, BJ,
international application was filed: I 93201757.7 18 June 1993 (18.06.93)	BEeta E	
(34) Countries for which the regional or		
	BE et a	
93870181.0 31 August 1993 (31.08.93) (34) Countries for which the regional or	E	With international search report.
	BE et a	
(71) Applicant (for all designated States except US): THE	PROC	
TER & GAMBLE COMPANY [US/US]; One Pr		
Gamble Plaza, Cincinnati, OH 45202 (US).		
(72) Inventors; and		
(75) Inventors/Applicants (for US only): GEBOES, Peter, Joannes [BE/BE]; Kapellestraat 230, B-2630 A		
	m meta	

(54) Title: CLEANING COMPOSITIONS WITH COMBINED HIGHLY HYDROPHILIC AND HIGHLY HYDROPHOBIC NONIONIC SURFACTANTS

#### (57) Abstract

The present invention encompasses cleaning compositions comprising a nonionic surfactant system containing a mixture of highly hydrophilic and highly hydrophobic nonionic surfactants. The present invention also encompasses a method of cleaning wherein said compositions are diluted in water before they are used to clean hard surfaces. The present invention also encompasses the diluted cleaning composition obtained in the method according to the present invention.

#### FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
ΑÜ	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	BU	Hungary	NO	Norway
BG	Bulgaria	<b>IE</b>	Ireland	NZ	New Zealand
BJ	Benin	П	Italy	PŁ	Poland
BR	Brazil ·	JP	Japan	PT	Portugal
BÝ	Belarus	KE	Kenya	RO	Romania
CA	Canada	KĠ	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic	SD	Sudan
CG	Congo		of Korea	SE	Sweden
CH	Switzerland	· KR	Republic of Korea	SI	Slovenia
CI	Côte d'Ivoire	KZ	Kazakhstan	SK	Slovaria
CM	Сапретооп	LI	Liechtenstein	SN	Senegal
CN	China	LK	Sri Lanka	TD	Chad
cs	Czechoslovakja	LU	Luxembourg	TG	Togo
CZ	Czech Republic	LV	Latvia	TJ	Tajikistan
DE	Germany	MC	Monaco	TT	Trinidad and Tobago
DK	Denmark	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	US	United States of America
FI	Finland	ML	Mali	UZ	Uzbekistan .
FR	France	MIN	Mongolia	VN	Vict Nam
GA	Gabon		-		

1

CLEANING COMPOSITIONS WITH COMBINED HIGHLY HYDROPHILIC AND HIGHLY HYDROPHOBIC NONIONIC SURFACTANTS

### Technical field

The present invention relates to cleaning compositions. Although the present invention applies primarily to hard surface cleaning compositions, it may also be of interest for other cleaning compositions including dishwashing and laundry detergent compositions.

#### Background

Compositions which can be used in diluted form are known in the art. In such compositions, the technical problem lies in sustaining upon dilution an efficient cleaning performance. Such compositions for dilute cleaning are described for instance in the European Patent EP-A- 0 503 219. In these compositions, the dilute cleaning performance is obtained by using alkanolamines. This '219 patent application also mentions the use of nonionic surfactants among which ethoxylated fatty alcohols ar listed.

Hard surface cleaning compositions comprising various surfactants, including highly ethoxylated nonionic surfactants, which can be used in diluted form are also known in the art. Representative of this prior art are the compositions described in European Patent Application EP 92870045.9.

There is a constant strive for the development of cleaning compositions with better performance in several respects. Better cleaning on all soils, including greasy soils and soap scum is of course a principal object, both for neat and diluted compositions which are often used in the cleaning of for instance hard surfaces. Such greasy soils are commonly brought by cooking with oils and fatty foods, and by simple contact with human skin, thus they are present on most surfaces, particularly in bathrooms and kitchens.

It is thus an object of the present invention to provide cleaning compositions which can be effectively used in diluted form, on various surfaces as well as in laundry applications, to clean various soils and stains. More specifically, it is a further object of the present invention to provide satisfactory shine performance and surface safety when used as hard surface cleaners.

It has now been found that this object can be efficiently met by formulating cleaning compositions comprising a specific nonionic surfactant system. Indeed, it has been found that cleaning compositions comprising at least a first nonionic surfactant with an HLB of at least 12(hydrophilic-lipophilic balance) and at least a second nonionic surfactant with an HLB below 10 and at least 4 less than the HLB of said first surfactant exhibit a great flexibility in the soils it may clean. These compositions unexpectedly exhibit outstanding cleaning performance upon dilution, compared to compositions comprising typical

nonionic surfactants while the cleaning performance of the neat, i.e. undiluted compositions is similar, regardless of the presence of said nonionic surfactant system.

In other words, it has now been found that the cleaning compositions comprising a nonionic surfactant system of at least a first surfactant with an HLB of at least 12 and at least a second surfactant with an HLB below 10 and at least 4 less than the HLB of said first surfactant exhibit outstanding cleaning performance when used both neat and diluted.

Specifically, the compositions according to the present invention are especially effective on greasy soils, both when used in hard water and also when used in soft water. Indeed, a particular advantage of the present invention is robustness to different usage conditions, e.g. variations in water hardness used for diluting the product.

This invention is also particularly useful as it allows to formulate compositions with reduced levels of other ingredients, yet retaining optimal cleaning performance, due to said combination of at least a first nonionic surfactant with an HLB of at least 12 and at least a second nonionic surfactant with an HLB below 10 and at least 4 less than the HLB of said first surfactant; also, the user needs less product to achieve the same task. This advantage is particularly valuable in terms of environmental compatibility.

#### Summary of the invention

The present invention encompasses an aqueous cleaning composition comprising from 0.5 % to 30 % by weight of the total composition of a nonionic surfactant system, whereby said nonionic surfactant system comprises:

- at least 0.1 % by weight of the total composition of a first surfactant with an HLB of at least 12;
- at least 0.1 % by weight of the total composition of a second surfactant with an HLB below 10 and at least 4 less than the HLB of said first surfactant.

The present invention encompasses both neat compositions as described hereinabove and diluted compositions. Diluted compositions are obtained by diluting in water the above neat compositions comprising from 0.5 % to 30 % by weight of the total composition of a nonionic surfactant system. Said diluted compositions comprise from 0.001 % to 5 % by weight of the total composition of said nonionic surfactant system.

The present invention also encompasses a method of cleaning hard surfaces wherein a composition comprising a nonionic surfactant system, whereby said nonionic surfactant system comprises at least 0.1 % by weight of the total composition of a first surfactant with an HLB of at least 12 and at least 0.1 % by weight of the total composition of a second surfactant with an HLB below 10 and at least 4 less than the HLB of said first surfactant, is diluted in water and subsequently applied to said hard surface.

#### Detailed description of the invention

The compositions according to the invention comprise, as an essential ingredient, a nonionic surfactant system comprising at least a nonionic surfactant with an HLB of at least 12, hereinafter referred to as highly hydrophilic surfactant and at least a nonionic surfactant with an HLB below 10 and at least 4 less than that of said highly hydrophilic surfactant, hereinafter referred to as highly hydrophobic surfactant.

ŧ

Suitable nonionic surfactants for the implementation of said surfactant system are alkoxylated alcohols or alkoxylated phenylalcohols which are commercially available with a variety of alcohol chain lengths and a variety of alkoxylation degrees. By simply varying the length of the chain of the alcohol and/or the degree of alkoxylation, alkoxylated alcohols or alkoxylated phenylalcohols can be obtained with different HLB values. It is to be understood to those ordinarily skilled in the art that the HLB value of any specific compound is available from the literature.

Suitable chemical processes for preparing the highly hydrophilic and highly hydrophobic nonionic surfactants for use herein include condensation of corresponding alcohols with alkylene oxide, in the desired proportions. Such processes are well known to the man skilled in the art and have been extensively described in the art. As an alternative, a great variety of alkoxylated alcohols suitable for use herein is commercially available from various suppliers.

The highly hydrophilic nonionic surfactants for use in the present invention have an HLB of at least 12, preferably above 14 and most preferably above 15. Those highly hydrophilic nonionic surfactants have been found to be particularly efficient for a rapid wetting of typical hard surfaces covered with greasy soils and to provide effective soil suspension.

The highly hydrophobic nonionic surfactants for use in the present invention have an HLB below 10, preferably below 9 and most preferably below 8.5. Those highly hydrophobic nonionic surfactants have been found to provide excellent grease cutting and emulsification properties.

Preferred highly hydrophilic nonionic surfactants to b used in the compositions according to the present invention

are surfactants having an HLB from 12 to 20 and being according to the formula  $RO-(C_2H_4O)_n(C_3H_6O)_mH$ , wherein R is a  $C_8$  to  $C_{22}$  alkyl chain or a  $C_8$  to  $C_{28}$  alkyl benzen chain, and wherein n+m is from 6 to 100 and n is from 0 to 100 and m is from 0 to 100, preferably n+m is from 21 to 50 and, n and m are from 0 to 50, and more preferably n+m is from 21 to 35 and, n and m are from 0 to 35. Throughout this description n and m refer to the average degree of the ethoxylation/propoxylation. The preferred R chains for use herein are the C8 to C22 alkyl chains. Examples of highly hydrophilic nonionic surfactants for use herein are LUTENSOL R A030 (HLB=17; R is a mixture of  $C_{13}$  and  $C_{15}$ alkyl chains, n is 30 and m is 0) commercially available from BASF, CETALOX R 50 (HLB= 18; R is a mixture of C16 and  $C_{18}$  alkyl chains, n is 50 and m is 0) commercially available from WITCO Alfonic R and 810-60 (HLB=12; R is a mixture of  $C_8$  and  $C_{10}$  alkyl chains, n is 6 and m is 0); and MARLIPAL R 013/400 (HLB= 18; R is a mixture of  $C_{12}$  and  $C_{14}$ , n is 40 and m is 0) commercially available from HULS.

Preferred highly hydrophobic nonionic surfactants to be used in the compositions according to the present invention are surfactants having an HLB of from 2 to 10 and being according to the formula RO-(C2H4O)n(C3H6O)mH, wherein R is a  $C_8$  to  $C_{22}$  alkyl chain or a  $C_8$  to  $C_{28}$  alkyl benzene chain, and wherein n+m is from 0.5 to 5 and n is from 0 to 5 and m is from 0 to 5, preferably n+m is from 0.5 to 4 and, n and m are from 0 to 4, more preferably n+m is from 1 to 4 and, n and m are from 0 to 4. The preferred R chains for use herein are the C8 to C22 alkyl chains. Examples of highly hydrophobic nonionic surfactants for use herein are DOBANOL R 91-2.5 (HLB= 8.1; R is a mixture of C9 and  $C_{11}$  alkyl chains, n is 2.5 and m is 0) commercially available from SHELL, LUTENSOL R AO3 (HLB=8; R is a mixture of C13 and C15 alkyl chains, n is 3 and m is 0) commercially available from BASF; Neodol 23-3 (HLB=7.9; R is a mixture of  $C_{12}$  and  $C_{13}$  alkyl chains, n is 3 and m is 0) and TERGITOL R 25L3

(HLB= 7.7; R is in the range of  $C_{12}$  to  $C_{15}$  alkyl chain length, n is 3 and m is 0) commercially available from UNION CARBIDE.

It is possible to use for each category of nonionic surfactants (highly hydrophilic or highly hydrophobic) either one of the nonionic surfactant belonging to said category or mixtures thereof.

The aqueous cleaning compositions according to the present invention comprise from 0.5% to 30% by weight of the total composition of said nonionic surfactant system, preferably from 2 % to 25 %, more preferably from 4 % to 20 %.

The compositions according to the present invention comprise said highly hydrophilic nonionic surfactant in an amount of at least 0.1 %, preferably of at least 0.5 %, more preferably of at least 2 %, and said highly hydrophobic nonionic surfactant in an amount of at least 0.1%, preferably of at least 0.5 %, more preferably of at least 2 %.

In the compositions according to the present invention, said highly hydrophilic and highly hydrophobic nonionic surfactants may be used in a weight ratio from one to another of from 0.1:1 to 1:0.1, preferably of from 0.2:1 to 1:0.2.

The present invention further encompasses diluted compositions which are obtainable by diluting the compositions described hereinabove, and said diluted compositions comprise from 0.001 % to 5 % by weight of the total composition of the nonionic surfactant mixture described hereinabove, preferably from 0.01 % to 0.5 %.

In a preferred embodiment of the present invention the compositions according to the present invention further

comprise an anionic surfactant. The prior art extensively describ s anionic surfactants, including alkyl benzene sulfates and sulfonates, alkyl ether sulfates, paraffin sulfonates, sulfonates of fatty acids and of fatty acid esters, sulpho succinates, sarcosinates, all of which are suitable for use herein. Said anionic surfactants can also be used in the form of their salts, including sodium, potassium, magnesium, ammonium and alkanol/alkyl ammonium salts. The preferred anionic surfactants for use herein are paraffin sulfonates or alkyl sulfates, preferably in the form of their sodium or ammonium salt. The aqueous cleaning compositions according to the present invention comprise up to 15% by weight of the total neat composition of anionic surfactant, preferably from 2 % to 6 %.

The cleaning compositions according to the present invention may further comprise additional surfactants, including other nonionic surfactants, amphoteric and zwitterionic surfactants and mixtures thereof. Typical of these are amine oxides, betaines, sulphobetaines and the like, which are well-known in the detergency art.

The compositions according to the present invention may further comprise an alkanolamine, or mixtures thereof, in amounts ranging from 0.1% to 10% by weight of the total neat composition, preferably from 0.5% to 5%, most preferably from 0.8% to 3%. At such levels, the alkanolamine has a buffering effect for alkaline products in the undiluted product, as well as an unexpected boosting effect on the cleaning performance of the diluted compositions. Suitable alkanolamines for use in the compositions according to the present include monoalkanolamines, dialkanolamines, trialkanolamines, alkylalkanolamines, dialkylalkanolamines and alkoxyalkanolamines. Preferred alkanolamines to be us d according to the present invention include monoethanolamine, triethanolamine, aminoethylpropanediol,

2-aminomethyl propanol, and ethoxyethanolamine. Particularly preferred are monoethanolamine, triethanolamine and ethoxyethanolamine.

The compositions according to the present invention may further comprise builders. Suitable builders for use herein include nitrilotriacetates (NTA), polycarboxylates, dipicolinates, citrates, water-soluble phosphates such as tri-polyphosphate and sodium ortho-and pyro-phosphates, silicates, ethylene diamine tetraacetate (EDTA), aminopolyphosphonates (DEQUEST), ether carboxylate builders such as in EP-A-286 167, phosphates, iminodiacetic acid derivatives such as described in EP-A-317 542, EP-262 112 and EP-A-399 133, and mixtures thereof. Preferred builders/sequestrants for use herein are selected from tetrapotassium pyrophosphate and citric acid. The aqueous cleaning compositions according to the present invention may comprise up to 15% by weight of the total neat composition of a builder, preferably from 1% to 10%.

The compositions according to the present invention may further comprise solvents. Suitable solvents for use herein include propylene glycol derivatives such as nbutoxypropanol or n- butoxypropoxypropanol, water-soluble CARBITOL R solvents or water-soluble CELLOSOLVE R solvents: water-soluble CARBITOL R solvents are compounds of the 2-(2-alkoxyethoxy) ethanol class wherein the alkoxy group is derived from ethyl, propyl or butyl; a preferred watersoluble carbitol is 2-(2-butoxyethoxy) ethanol also known as butyl carbitol. Water-soluble CELLOSOLVE R solvents are compounds of the 2-alkoxyethoxy ethanol class, with 2butoxyethoxyethanol being preferred. Other suitable solvents include benzyl alcohol, and diols such as 2-ethyl-1, 3-hexanediol and 2,2,4-trimethyl-1,3-pentanediol and mixtures thereof. Preferred solvents for use herein are nbutoxypropoxypropanol, BUTYL CARBITOL R and mixtures The compositions according to the present

invention comprise up to 15% by weight of the total neat composition of a solvent, preferably from 3% to 10%.

The compositions according to the present invention may also comprise minor ingredients to provide aesthetic or additional product performance benefits. Typical minor ingredients include perfumes, dyes, optical brighteners, soil suspending agents, detersive enzymes, gel-control agents, thickeners, freeze-thaw stabilizers, bactericides, preservatives, hydrotopes like sodium cumene sulfonate and the like.

The present invention further encompasses a method for the cleaning of hard surfaces. According to the present invention, an aqueous cleaning composition comprising from 0.5 % to 30 % by weight of the total composition of a nonionic surfactant system, whereby said nonionic surfactant system comprises at least 0.1 % by weight of the total composition of a first surfactant with an HLB of at least 12 and at least 0.1 % by weight of the total composition of a second surfactant with an HLB below 10 and at least 4 less than said first surfactant's HLB, is diluted in water so as to obtain from 0.001 % to 5 % by weight of the total composition of said nonionic surfactant system and subsequently applied to said hard surface.

In the method according to the present invention, said composition is diluted into 10 to 200 times its weight of water, more preferably 50 to 150 times, before it is applied to said hard surface.

The present invention is further illustrated by the following examples.

#### Example I

The following compositions were tested for their cl aning performance when used diluted on greasy soil.

The following compositions were made by mixing the listed ingredients in the listed proportions:

Compositions:	#1	#2	#3
Ingredients:		We	ight %
Sodium paraffin sulfonate	3	3	3
Alcohol ethoxylate 7EO (1)	4	-	-
Alcohol ethoxylate 30EO (2)		. 3	2
Alcohol ethoxylate 3EO (3)	-	1	2
Sodium Citrate	3	. 3	· 3
Butylcarbitol R	4	4	4
Triethanolamine	1	1	1
water & minors	1	up to	100%

These compositions were evaluated for their grease removal efficiency. A high index is correlated to a high performance.

Dilute cleaning performance (index)

Compositions	#1	#2	#3
Hard water	100	370	350
Soft water	100	200	220

In the examples hereinabove, (1) is an ethoxylate nonionic surfactant representative of the prior art with an HLB of 12. Compounds (2) and (3) are ethoxylate nonionic surfactants according to the present invention. (2) is a highly hydrophilic nonionic surfactant with an HLB of 17, wherein R is a mixture of C13 and C15 alkyl chains and n is

30. (3) is a highly hydrophobic nonionic surfactant with an HLB of 8, wherein R is a mixture of C13 and C15 alkyl chains and alkyl chains and n is 3.

Compositions 1-3 described hereinabove can be used neat or diluted. In the present cleaning performance test these compositions are diluted in 64 times their weight of water and subsequently applied to a hard surface.

Typical dilute grease cleaning results obtained for such compositions are given for the examples hereinabove. Composition #1 which comprises a nonionic surfactant with an moderate HLB (12) gets a cleaning index of 100 and is used as a reference.

According to the present invention, the compositions #2 and #3 which comprise a nonionic surfactant system comprising alcohol ethoxylate 30EO (an highly hydrophilic nonionic surfactant) and alcohol ethoxylate 3EO (an highly hydrophobic nonionic surfactant) exhibit outstanding cleaning performance upon dilution both in hard and soft water. Indeed, when used with hard water the compositions #2 and #3 are almost four times more efficient than the compositions #2 and #3 are two times more efficient than the compositions #2 and #3 are two times more efficient than the composition #1.

The above results surprisingly show that the use of diluted compositions according to the present invention provides a significant benefit in grease cleaning compared to the composition comprising only nonionic surfactant with a moderate HLB (for example 12), both in hard and soft water.

## Example II

A concentrated composition is prepared with the following composition:

Ingredients	Weight %
Alfonic 810-60 (C <sub>8</sub> -C <sub>10</sub> E <sub>6</sub> )	20.0
Neodol 23-3 $(C_{12}-C_{13} E_3)$	4.0
Coconut fatty acid	0.6
Monoethanolamine	1.0
Sodium cumene sulfonate	1.5
Minors and water	Balance

This composition provides good performance while requiring less packaging.

#### What is claimed is:

- 1. An aqueous cleaning composition comprising from 0.5 % to 30 % by weight of the total composition of a nonionic surfactant system, characterized in that said nonionic surfactant system comprises:
  - at least 0.1 % by weight of the total composition of a first highly hydrophilic surfactant with an HLB of at least 12;
  - at least 0.1 % by weight of the total composition of a second highly hydrophobic surfactant with an HLB below 10 and at least 4 less than the HLB of said first surfactant.
- 2. A composition according to claim 1 wherein in said nonionic surfactant system said highly hydrophilic surfactants have an HLB above 14, preferably above 15, and said highly hydrophobic surfactants have an HLB below 9, preferably below 8.5.
- 3. A composition according to any of the preceding claims wherein the amount of said highly hydrophilic nonionic surfactant in the total composition is at least 0.5 % by weight, preferably at least 2 %, and the amount of said highly hydrophobic nonionic surfactant in the total composition is at least 0.5 % by weight, preferably at least 2 %.
- 4. A composition according to any of the preceding claims wherein the weight ratio of said highly hydrophilic nonionic surfactant to said highly hydrophobic nonionic surfactant is from 0.1:1 to 1:0.1, preferably from 0.2:1 to 1:0.2.

- wherein the highly hydrophilic nonionic surfactant is a nonionic of the formula RO-(C<sub>2</sub>H<sub>4</sub>)<sub>n</sub>(C<sub>3</sub>H<sub>4</sub>)<sub>m</sub>H wherein R is a C<sub>8</sub> to C<sub>22</sub> alkyl chain or a C<sub>8</sub> to C<sub>28</sub> alkyl benzene chain, and wherein n+m is from 6 to 100 and n and m are from 0 to 100, preferably n+m is from 21 to 50 and n and m are from 0 to 50, most preferably n+m is from 21 to 35 and n and m are from 0 to 35.
- 6. A composition according to any of the preceding claims wherein the highly hydrophobic nonionic surfactant is a nonionic of the formula RO-(C<sub>2</sub>H<sub>4</sub>)<sub>n</sub>(C<sub>3</sub>H<sub>4</sub>)<sub>m</sub>H wherein R is a C<sub>8</sub> to C<sub>22</sub> alkyl chain or a C<sub>8</sub> to C<sub>28</sub> alkyl benzene chain, and wherein n+m is from 0.5 to 5 and n and m are from 0 to 5, preferably n+m is from 0.5 to 4 and n and m are from 0 to 4, and most preferably n+m is from 1 to 4 and n and m are from 0 to 4.
- 7. A composition according to any of the preceding claims which further comprises up to 15% by weight of the total composition of an anionic surfactant.
- 8. A composition according to claim 7, wherein said anionic surfactant is a paraffin sulfonate in the form of its sodium salt.
- 9. A method for the cleaning of a hard surface wherein a cleaning composition according to any of the preceding claims is diluted in water and subsequently applied to said hard surface.
- 10. A method according to claim 9, wherein said composition is diluted into 10 to 200 times its weight of water, more preferably 50 to 150 times, before it is subsequently applied to said hard surface.

11. A diluted hard surface cleaning composition obtainable by diluting in water a composition according to claims 1 to 8, comprising from 0.001 % to 5 %, preferably from 0.01 % to 0.5 % by weight of the total composition of said nonionic surfactants.

#### INTERNATIONAL SEARCH REPORT

Inte . .ional application No.

PCT/US93/11293

A. CLASSIFICATION OF SUBJECT MATTER  IPC(5) :C11D 1/66, 1/72, 1/722, 1/825.				
US CL:252/173, 174.21, 174.22, DIG. 1, DIG. 14. According to International Patent Classification (IPC) or to both national classification and IPC				
	DS SEARCHED			
	ocumentation searched (classification system followed	by classification symbols)	<del></del>	
	252/173, 174.21, 174.22, DIG. 1, DIG. 14.			
Documentat	ion searched other than minimum documentation to the	extent that such documents are included	in the fields searched	
· .				
Electronic o	data base consulted during the international search (na	me of data base and, where practicable	search terms used)	
NONE				
C. DOC	UMENTS CONSIDERED TO BE RELEVANT			
Category*	Citation of document, with indication, where ap	propriate, of the relevant passages	Relevant to claim No.	
X	US,A, 5,145,850 (DEGUCHI ET A	L.) 13 October 1992 see	1-3	
	col. 2, lines 20-46, especially line			
Υ	21; Table 1, Example Composition	rs 2, 5 & 6.	1-3	
		·		
			·	
	·			
	·	•		
Furt	ner documents are listed in the continuation of Box C	. See patent family annex.		
*A* do	ecial categories of cited documents: cument defining the general state of the art which is not considered	"T" later document published after the int date and not in conflict with the applic principle or theory underlying the inv	ation but cited to understand the	
	be part of particular relevance rlier document published on or after the international filing date	"X" document of particular relevance; the	e claimed invention cannot be	
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other				
.O. 90	ecial reason (as specified)  cument referring to an oral disclosure, use, exhibition or other cans	considered to involve an inventive combined with one or more other suc being obvious to a person skilled in t	step when the document is high documents, such combination	
*P* document published prior to the international filing date but later than *&* document member of the same patent family the priority date claimed				
Date of the actual completion of the international search  15 January 1994  Date of mailing of the international search 17 FEB 1994				
Name and	Name and mailing address of the ISA/US  Authorized officer  Authorized officer			
Box PCT	Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231			
l	Talanhara No. (702) 209-2520			